



FavorPrep™ miRNA Isolation Kit

Cat.: FAMIK000
 FAMIK001
 FAMIK002
(For Research Use Only)

Introduction:

FavorPrep™ miRNA Isolation Kit is designed for purification of microRNAs (miRNAs) and other small cellular RNAs from tissue samples or cultured cells. Purification of miRNAs allows research into biological significant pathways for gene regulation.

The standard protocols for isolating total RNA and mRNA are not optimized for isolation of small RNA molecules and result in the loss of substantial amounts of miRNAs and other small RNA. In addition, removal of the predominant larger RNAs is required for accurate analysis of miRNA expression by qPCR or microarray analysis.

This kit is specifically designed for purification of small RNA with minimal contamination from large RNA molecules or genomic DNA. The method employs a spin column with a silica-based fiber matrix that binds RNA in the presence of a chaotropic salt. The method is based on the selective binding of RNA molecules of different sizes to the silica-based fiber matrix when different ethanol concentrations are present in the solvent.

Kit Contents:

Cat. No:	FAMIK000 (5 preps)	FAMIK001 (100 preps)	FAMIK002 (50 preps)
Lysis Buffer	1.5 ml	25 ml	12 ml
2M NaOAc, pH 5.2	150 µl	2.5 ml	1.2 ml
Wash Buffer 2 (concentrate) ^a	0.3 ml	5 ml	3 ml
Release Buffer	0.32 ml	5.5 ml	2.8 ml
RNA Column	10 pcs	200 pcs	100 pcs
Collection Tube	10 pcs	200 pcs	100 pcs
Preparation of Wash Buffer 2 by adding ethanol (96 ~ 100%) and Store at RT.			
Ethanol volume for Wash Buffer 2 ^a	1.2 ml	20 ml	12 ml

Specification:

Principle:	mini spin column (silica matrix)
Sample size:	up to 1 x10 ⁶ cultured cells up to 100 mg tissue
Operation time:	~ 30 minutes
Column applicability:	centrifugation and vacuum

Additional requirement to be provided by user

1. Microcentrifuge capable of speed at ~12,000 rpm
2. 1.5 ml microcentrifuge tube
3. 96~100 % ethanol
4. Water-Saturated Phenol
5. Chloroform
6. Vortex
7. Water bath or dry bath

Important Notes:

1. Buffers provided in this system contain irritants. Wear gloves and lab coat when handling these buffers.
2. Store the kit at room temperature.
3. Caution: phenol and chloroform are hazardous to human health. perform the procedures involving phenol and chloroform in a chemical fume hood.
4. Add required volume of ethanol (96- 100 %) to Wash Buffer 2 when first open. **Store the solution at room temperature.**

Protocol:

Read the Important Note before starting the following steps.

HINT: Preheat Release Buffer to 65°C for step 15.

1. Add 200 µl Lysis Buffer into the tube containing up to 100 mg tissue or 1x10⁶ cultured cell pellet.
2. Vigorous mixing by vortexing. Incubate at room temperature for 10 minutes.
3. Add 20 µl 2M NaOAc, pH 5.2.
4. Add 180 µl water-saturated phenol and 40 µl chloroform into the tube, vortex vigorously for 2 minutes.
5. Centrifuge at 12,000 rpm for 3 minutes. Transfer the upper phase into a new 1.5 ml centrifuge tube.
6. Add a 35% volume of absolute ethanol to the upper phase and mix well by shaking vigorously.
-- If the upper phase volume is 200 µl, add 108 µl of absolute ethanol to upper phase.
7. Place a RNA Column in a Collection Tube and transfer the ethanol-added mixture to the RNA Column. Incubate for 1 minute.
8. Centrifuge at 12,000 rpm for 30 seconds (Large RNA bound to the column membrane). Collect the filtrate and transfer the filtrate to a new 1.5 ml microcentrifuge tube.
9. Add a 70% volume of absolute ethanol to the filtrate and mix well by shaking vigorously.
-- If the filtrate (ethanol-contained, 108 µl) volume is 308 µl, add 338 µl of absolute ethanol to the filtrate. The final concentration ethanol of whole mixture is 70%.
10. Place a new RNA Column in a Collection Tube then transfer the mixture to the RNA Column. Incubate for 1 minute.
11. Centrifuge at 12,000 rpm for 30 seconds (miRNA bound to the column membrane).
12. Add 200 µl Wash Buffer 2 (ethanol added). Incubate for 1 minute.
13. Centrifuge at 12,000 rpm for 1 minute to completely remove the residue liquid.
14. Put the RNA Column to a new 1.5 ml centrifuge tube.
15. Add 50 µl Release Buffer (preheated to 65°C) to the center of column. Incubate for 3 minutes.
16. Centrifuge at 12,000 rpm for 3 minute to recover miRNA. (Note: The purified miRNA can be further concentrated by a standard ethanol precipitation procedure and then re-dissolved in a small volume ddH₂O or TE, pH 8.0).
17. Use 1/5 volume to run on a mini agarose gel (or more accurately, a polyacrylamide gel) to check its quality. The majority of RNA visible on the gel should be <100 nt in size, with the major bands corresponding to tRNAs. The 5S and 5.8S rRNA species may also be visible. These tRNA and small rRNA bands should be clear and distinct. miRNA (21-22 nt) are typically not visible on the gel image.